

Fibroadenoma

This leaflet tells you about fibroadenoma. It explains what a fibroadenoma is, how it's diagnosed and what will happen if it needs to be followed up or treated.



Benign breast conditions information
provided by Breast Cancer Care



Breast Cancer Care doesn't just support people when they've been diagnosed with breast cancer

We also highlight the importance of early detection and provide up-to-date, expert information on breast conditions and breast health.

If you have a question about breast health or breast cancer you can call us free on **0808 800 6000** or visit **breastcancercare.org.uk**

We hope you found this information useful.
If you'd like to help ensure we're there for other people when they need us visit **breastcancercare.org.uk/donate**

Central Office

Breast Cancer Care

Chester House

1-3 Brixton Road

London SW9 6DE


Phone: 0345 092 0800

Email: info@breastcancercare.org.uk

What is a fibroadenoma?

A fibroadenoma is a very common benign (not cancer) breast condition.

The breast



Fibroadenomas often develop during puberty so are mostly found in young women, but they can occur in women of any age. Men can also get fibroadenomas, but this is very rare.

Symptoms of fibroadenoma

A fibroadenoma is usually felt as a lump in the breast which is smooth to the touch and moves easily under the skin.

Fibroadenomas are usually painless, but sometimes they may feel tender or even painful, particularly just before a period.

Types of fibroadenoma

Simple fibroadenoma

Most fibroadenomas are about 1–3cm in size and are called simple fibroadenomas. When looked at under a microscope, simple fibroadenomas will look the same all over.

Simple fibroadenomas do not increase the risk of developing breast cancer in the future.

Complex fibroadenoma

Some fibroadenomas are called complex fibroadenomas. When these are looked at under a microscope, some of the cells have different features.

Having a complex fibroadenoma can very slightly increase the risk of developing breast cancer in the future.

Giant or juvenile fibroadenoma

Occasionally, a fibroadenoma can grow to more than 5cm and may be called a giant fibroadenoma. Those found in teenage girls may be called juvenile fibroadenomas.

Causes of fibroadenoma

It's not known what causes a fibroadenoma. It's thought that it probably occurs because of increased sensitivity to the hormone oestrogen.

Breasts are made up of lobules (milk-producing glands) and ducts (tubes that carry milk to the nipple). These are surrounded by glandular, fibrous and fatty tissue. This tissue gives breasts their size and shape.

Fibroadenomas develop from a lobule. The glandular tissue and ducts grow over the lobule and form a solid lump.

Diagnosis

If you find a breast lump, see your GP. They're likely to refer you to a breast clinic where you'll be seen by specialist doctors or nurses.

At the clinic you'll have a range of tests. These may include:

- A mammogram (breast x-ray)
- An ultrasound scan (using sound waves to produce an image)
- A core biopsy (using a hollow needle to take a sample of breast tissue to be looked at under a microscope – several tissue samples may be taken at the same time)
- A fine needle aspiration (FNA) (using a fine needle and syringe to take a sample of cells to be looked at under a microscope)

Fibroadenomas are often easier to identify in younger women. If you're in your early 20s or younger, your fibroadenoma may be diagnosed with a breast examination and ultrasound only. However, if there's any uncertainty about the diagnosis, a core biopsy or FNA will be done.

If you're under 40, you're more likely to have an ultrasound than a mammogram. Younger women's breast tissue can be dense which can make the x-ray image less clear so normal changes or benign breast conditions can be harder to identify. However, for some women under 40, mammograms may still be needed to complete the assessment.

You can call us free on **0808 800 6000** if you'd like more information about any tests you're having, or see our booklet **Your breast clinic appointment**.

Treatment and follow-up

In most cases you'll not need any treatment or follow-up if you have a fibroadenoma. Usually you'll only be asked to go back to your GP or the breast clinic if it gets bigger or you notice a change.

Most fibroadenomas stay the same size. Some get smaller and some eventually disappear over time. A small number of fibroadenomas get bigger, particularly those in teenage girls. Fibroadenomas can also get bigger during pregnancy and breastfeeding or while taking hormone replacement therapy (HRT), but usually reduce in size again afterwards.

Surgery

Sometimes an operation, called an excision biopsy, is needed to remove a fibroadenoma if it's a large, complex or juvenile fibroadenoma. You can also ask to have a fibroadenoma removed. This is usually performed under general anaesthetic.

Your surgeon may use dissolvable stitches placed under the skin which will not need to be removed. However, if a non-dissolvable stitch is used, they'll need to be taken out about a week after surgery. You'll be given information about looking after the wound before you leave hospital.

Vacuum assisted (VAC/VAB) excision biopsy

You may be offered a vacuum assisted excision biopsy to remove the fibroadenoma. This is a way of removing small fibroadenomas under local anaesthetic, without having an operation under general anaesthetic.

After an injection of local anaesthetic, a small cut is made in the skin. A hollow probe connected to a vacuum device is placed through this. Using an ultrasound as a guide, the fibroadenoma is sucked through the probe by the vacuum into a collecting chamber. The biopsy device is used in this way until all of the fibroadenoma has been removed. This may mean that surgery can be avoided.

The removed tissue is sent to a laboratory and examined under a microscope.

This procedure can cause some bruising and pain for a few days afterwards. Removing a fibroadenoma does not usually affect the shape of the breast, but there may be a slight dent in the breast where it has been removed.

Staying breast aware

For most women, having a fibroadenoma does not increase the risk of developing breast cancer.

If you were diagnosed with a complex fibroadenoma, you may be worried that your risk is very slightly increased. However, this does not necessarily mean you'll develop breast cancer in the future.

It's important to continue to be breast aware and go back to your GP if you notice any changes in your breasts, regardless of how soon these occur after your diagnosis of a fibroadenoma.

You can find out more about being breast aware in our booklet **Know your breasts: a guide to breast awareness and screening**.

About this leaflet

Fibroadenoma was written by Breast Cancer Care's clinical specialists, and reviewed by healthcare professionals and people affected by breast problems.



For a full list of the sources we used to research it:

Phone **0345 092 0808**

Email **publications@breastcancercare.org.uk**



You can order or download more copies from

breastcancercare.org.uk/publications



We welcome your feedback on this publication:

breastcancercare.org.uk/feedback



For a large print, Braille, or audio CD version:

Phone **0345 092 0808**

Email **publications@breastcancercare.org.uk**